

## REFERENCES

### Chapter 69

1. Mayson KV, Beestra JE, Choi PT. The incidence of postoperative complications in the PACU. *Can J Anesth*. 2005;52:A62.
2. Hines R, Barash PG, Watrous G, O'Connor T. Complications occurring in the postanesthesia care unit: a survey. *Anesth Analg*. 1992 Apr;74(4):503–509. PMID 1554116
3. Apfel CC, Läärä E, Koivuranta M, Greim CA, Roewer N. A simplified risk score for predicting postoperative nausea and vomiting: conclusions from cross-validations between two centers. *Anesthesiology*. 1999 Sep;91(3):693–700. PMID 10485781
4. Apfel CC, Korttila K, Abdalla M, et al. A factorial trial of six interventions for the prevention of postoperative nausea and vomiting. *N Engl J Med*. 2004 Jun;350(24):2441–2451. PMID 15190136
5. Eikermann M, Blobner M, Groeben H, et al. Postoperative upper airway obstruction after recovery of the train of four ratio of the adductor pollicis muscle from neuromuscular blockade. *Anesth Analg*. 2006 Mar;102:937–942. PMID 16492855
6. American Society of Anesthesiologists Task Force on Postanesthetic Care. Practice guidelines for postanesthetic care: a report by the American Society of Anesthesiologists Task Force on Postanesthetic Care. *Anesthesiology*. 2002 Mar;96:742–752. PMID 11873052
7. Gan TJ, Meyer TA, Apfel CC, et al. Society for Ambulatory Anesthesia guidelines for the management of postoperative nausea and vomiting. *Anesth Analg*. 2007 Dec;105(6):1615–1628. PMID 18042859