

## REFERENCES

### Chapter 82

1. Choukalas CG, Walter J, Glick D, Connor MF, Tung A, Dinwiddie SH, Nunnally ME. Mask ventilation, hypocapnia, and seizure duration in electroconvulsive therapy. *J. Clin. Anesth.* 2010 Sep;22(6):415–419. PMID 2086961
2. Eser D, Nothdurfter C, Schüle C, et al. The influence of anaesthetic medication on safety, tolerability and clinical effectiveness of electroconvulsive therapy. *World J. Biol. Psychiatry.* 2010 Mar;11(2 pt 2):447–456. PMID 19462341
3. Hooten WM, Rasmussen KG Jr. Effects of general anesthetic agents in adults receiving electroconvulsive therapy: a systematic review. *J. ECT.* 2008 Sep;24(3):208–223. PMID 18628717
4. Kadar AG, Ing CH, White PF, Wakefield CA, Kramer BA, Clark K. Anesthesia for electroconvulsive therapy in obese patients. *Anesth. Analg.* 2002 Feb;94(2):360–361. PMID 11812699
5. Tan HL, Lee CY. Comparison between the effects of propofol and etomidate on motor and electroencephalogram seizure duration during electroconvulsive therapy. *Anaesth. Intensive Care.* 2009 Sep;37(5):807–814. PMID 19775046
6. Krystal AD, Watts BV, Weiner RD, Moore S, Steffens DC, Lindahl V. The use of flumazenil in the anxious and benzodiazepine-dependent ECT patient. *J. ECT.* 1998 Mar;14(1):5–14. PMID 9661088