

REFERENCES

Chapter 84

1. Shapiro F. Anesthesia for outpatient cosmetic surgery. *Current Opinion in Anaesthesiology*. 2008 Dec;21(6):704–710. PMID 18997521
2. Iverson R, ASPS Task Force on Patient Safety in Office-Based Surgery Facilities: Patient Safety in office-based surgery facilities: I. Procedures in the office-based surgery setting. *Plast. Reconstr. Surg.* 2002 Oct;110(5):1337–1342. PMID 12360080
3. Rohrich RJ, Beran SJ. Is liposuction safe? *Plast. Reconstr. Surg.* 1999 Sep;104(3):819–22. PMID 10456537
4. Grazer FM, de Jong RH. Fatal outcomes from liposuction: Census survey of cosmetic surgeons. *Plast. Reconstr. Surg.* 2000 Jan;105(1):436–446. PMID 10627013
5. Culliford AT, Spector JA, Flores RL, et al. Intraoperative sensorcaine significantly improves postoperative pain management in outpatient reduction mammoplasty. *Plast. Reconstr. Surg.* 2007 Sep;120(4):840–844. PMID 17805109
6. Taghinia AH, Shapiro FE, Slavin SA. Dexmedetomidine in aesthetic facial surgery: improving anesthetic safety and efficacy. *Plast. Reconstr. Surg.* 2008;121:269–271. PMID 18176230
7. De Jong RH, et al. Perioperative Management of Cosmetic Liposuction. *Plast. Reconstr. Surg.* 2001;107(4):1039–1041. PMID 11252101
8. Klein JA. Tumescence technique for regional anesthesia permits lidocaine doses of 35 mg/kg. for liposuction. *J. Dermatol. Surg. Oncol.* 1990 Mar;16:248–263. PMID 2179348
9. Lelli GJ, Lisman RD. Blepharoplasty complications. *Plast. Reconstr. Surg.* 2010 Mar;125(3):1007–1017. PMID 20195127