

REFERENCES

Chapter 120

1. Baumann A, Audibert G, Klein O. Continuous intravenous lidocaine in the treatment of paralytic ileus due to severe spinal cord injury. *Acta Anaesthesiol Scand.* 2009 Jan;53:128–130. PMID 18945248
2. Borgeat A, Aguirre J. Update on local anesthetics. *Curr Opin Anaesthesiol.* 2010 Aug;23:466–471. PMID 20610984
3. Forster JG, Rosenberg PH. Clinically useful adjuvants in regional anesthesia. *Curr Opin Anaesthesiol.* 2003 Oct;16:477–478. PMID 17021500
4. Groudine SB, Fisher HA. Intravenous lidocaine speeds return of bowel function, decreases postoperative pain and shortens hospital stay in patients. *Anesth Analg.* 1998 Feb;86(2):235–239. PMID 9459225
5. Moiniche S, Mikkelsen S. et. al. A qualitative systematic review of incisional local anesthesia for postoperative pain relief after abdominal operations. *Br J Anaesth.* 1998 Sep;81(3):377–383. PMID 9861124
6. Sinnott CJ, Cogswell LP II et. al. I. On the mechanism by which epinephrine potentiates lidocaine's peripheral nerve block. *Anesthesiology.* 2003 Jan;98(1):181–188. PMID 12502995
7. Thorton PC, Grant SA, Breslin DS. Adjuncts to local anesthetics in peripheral nerve blockade. *Int Anesthesiol Clin.* 2010 Fall;48:59–70. PMID 20881527
8. Visser L. Epidural anaesthesia. *Update Anaesth.* 2001;(13) [article 11].
9. Warren DT, Liu SS. Neuraxial anesthesia. In: Longnecker DE, Brown DL, Newman F, Zapol M, eds. *Anesthesiology.* New York: McGraw-Hill; 2008.