

REFERENCES

Chapter 123

1. Motiani P, Chaudhary S, Bahl N, Sethi AK. Intrathecal sufentanil versus fentanyl for lower limb surgeries—a randomized controlled trial. *J Anaesthesiol Clin Pharmacol.* 2010 Jan;26(4):507–513. PMID 21804710
2. Sakura S, Sumi M, Sakaguchi Y, Saito Y, Kosaka Y, Drasner K. The addition of phenylephrine contributes to the development of transient neurologic symptoms after spinal anesthesia with 0.5% tetracaine. *Anesthesiology.* 1997 Oct;87(4):771–778. PMID 9357877
3. Zayas VM, Liguori GA, Chisholm MF, Susman MH, Gordon MA. Dose response relationships for isobaric spinal mepivacaine using the combined spinal epidural technique. *Anesth Analg.* 1999 Nov;89(5):1167–1171. PMID 10553828
4. McNamee DA, McClelland AM, Scott S, Milligan KR, Westman L, Gustafsson U. Spinal anaesthesia: comparison of plain ropivacaine 5 mg ml⁻¹ with bupivacaine 5 mg ml⁻¹ for major orthopaedic surgery. *Br J Anaesth.* 2002 Nov;89(5):702–706. PMID 12393766
5. Panni MK, George RB, Allen TK, et al. Minimum effective dose of spinal ropivacaine with and without fentanyl for postpartum tubal ligation. *Int J Obstet Anesth.* 2010 Oct;19(4):390–394 [Epub September 15, 2010]. PMID 20832280
6. Casati A, Danelli G, Berti M, et al. Intrathecal 2-chloroprocaine for lower limb outpatient surgery: a prospective, randomized, double-blind, clinical evaluation. *Anesth Analg.* 2006;103(1):234–238 [table of contents]. PMID 16790659
7. Liu SS. Optimizing spinal anesthesia for ambulatory surgery. *Reg Anesth.* 1997;22(6):500–510. PMID 9425964
8. Smith KN, Kopacz DJ, McDonald SB. Spinal 2-chloroprocaine: a dose-ranging study and the effect of added epinephrine. *Anesth Analg.* 2004 Jan;98(1):81–88 [table of contents]. PMID 14693591