

REFERENCES

Chapter 128

1. Frederickson MJ, Ball CM, Dalglish AJ. Analgesic effectiveness of a continuous versus single-injection interscalene block for minor arthroscopic shoulder surgery. *Reg Anesth Pain Med.* 2010 Jan-Feb;35(1):28–33. PMID 20048655
2. Ilfeld BM. Continuous peripheral nerve blocks: a review of the published evidence. *Anesth Analg.* 2011 Oct;113(4):904–925. PMID 21821511
3. Boezaart AP. Perineural infusion of local anesthetics. *Anesthesiology.* 2006 Apr;104(4):872–880. PMID 16571984
4. Wang AZ, Gu L, Zhou QH, Ni WZ, Jiang W. Ultrasound-guided continuous femoral nerve block for analgesia after total knee arthroplasty: catheter perpendicular to the nerve versus catheter parallel to the nerve. *Reg Anesth Pain Med.* 2010 Mar-Apr;35(2):127–131. PMID 20301819
5. Morin AM, Kranke P, Wulf H, Stienstra R, Eberhart LH. The effect of stimulating versus nonstimulating catheter techniques for continuous regional anesthesia: a semiquantitative systematic review. *Reg Anesth Pain Med.* 2010 Mar-Apr;35(2):194–199. PMID 20216037
6. Mariano ER, Loland VJ, Bellars RH, et al. Ultrasound guidance versus electrical stimulation for infraclavicular brachial plexus perineural catheter insertion. *J Ultrasound Med.* 2009 Sep;28(9):1211–1218. PMID 19710219
7. Mariano ER, Cheng GS, Choy LP, et al. Electrical stimulation versus ultrasound guidance for popliteal-sciatic perineural catheter insertion: a randomized controlled trial. *Reg Anesth Pain Med.* 2009 Sep-Oct;34(5):480–485. PMID 19920423
8. Fredrickson MJ, Ball CM, Dalglish AJ. A prospective randomized comparison of ultrasound guidance versus neurostimulation for interscalene catheter placement. *Reg Anesth Pain Med.* 2009 Nov-Dec;34:590–594. PMID 19916253
9. Ilfeld BM. Continuous peripheral nerve blocks in the hospital and at home. *Anesthesiol Clin.* 2011 Jun;29(2):193–211. PMID 21620338
10. Salinas FV, Liu SS, Mulroy MF. The effect of single-injection femoral nerve block versus continuous femoral nerve block after total knee arthroplasty on hospital length of stay and long-term functional recovery within an established clinical pathway. *Anesth Analg.* 2006 Apr;102(4):1234–1239. PMID 16551930
11. Horlocker TT, Wedel DJ, Benzoni H, et al. Regional anesthesia in the anticoagulated patient: defining the risks (the second ASRA consensus conference on neuraxial anesthesia and anticoagulation). *Reg Anesth Pain Med.* 2003 May-Jun;28:172–197. PMID 12772135
12. Horlocker TT, Wedel DJ, Rowlingson JC, et al. Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American Society of Regional Anesthesia and Pain Medicine evidence-based guidelines (3rd ed.). *Reg Anesth Pain Med.* 2010 Jan-Feb;35:64–101. PMID 20052816
13. Bergman BD, Hebl JR, Kent J, et al. Neurologic complications of 405 consecutive continuous axillary catheters. *Anesth Analg.* 2003 Jan;96(1):247–252. PMID 12505961
14. Bleckner LL, Bina S, Kwon KH, et al. Serum ropivacaine concentrations and systemic local anesthetic toxicity in trauma patients receiving long-term continuous peripheral nerve block catheters. *Anesth Analg.* 2010 Feb;110(2):630–634. PMID 19955504