

## REFERENCES

### Chapter 149

1. Eason MJ, Wyatt R. Paravertebral thoracic block—a reappraisal. *Anaesthesia*. 1979 Jul-Aug;34(7):638–642. PMID 517716
2. Karmakar MK. Thoracic paravertebral block. *Anesthesiology*. 2001 Sep;95(3):771–780. PMID 11575553
3. Cheema S, Richardson J, McGurgan P. Factors affecting the spread of bupivacaine in the adult thoracic paravertebral space. *Anaesthesia*. 2003 Jul;58(7):684–687. PMID 12886912
4. Naja Z, Lonnqvist PA. Somatic paravertebral nerve blockade. Incidence of failed block and complications. *Anaesthesia*. 2001 Dec;56(12):1184–1188. PMID 11736777
5. Karmakar MK, Ho AM, Law BK, Wong ASY, Shafer SL, Gin T. Arterial and venous pharmacokinetics of ropivacaine with and without epinephrine after thoracic paravertebral block. *Anesthesiology*. 2005;103(4):704–711. PMID 16192762
6. Richardson J, Sabanathan S, Jones J, Shah RD, Cheema S, Mearns AJ. A prospective, randomized comparison of preoperative and continuous balanced epidural or paravertebral bupivacaine on post-thoracotomy pain, pulmonary function and stress responses. *Br J Anaesth*. 1999 Sep;83(3):387–392. PMID 10655907
7. Ben-Ari A, Moreno M, Chelly JE, Bigeleisen PE. Ultrasound-guided paravertebral block using an intercostal approach. *Anesth Analg*. 2009 Nov;109(5):1691–1694. PMID 19843811
8. Cowie B, McGlade D, Ivanusic J, Barrington MJ. Ultrasound-guided thoracic paravertebral blockade: a cadaveric study. *Anesth Analg*. 2010 Jun;110(6):1735–1739. PMID 20435949