

REFERENCES

Chapter 163

1. Buckenmaier CC, Shields CH, Auton AA, et al. Continuous peripheral nerve block in combat casualties receiving low-molecular weight heparin. *Br J Anaesth*. 2006 Dec;97(6):874–877. PMID 17032662
2. Capdevila X, Bringuier S, Borgeat A. Infectious risks of continuous peripheral nerve blocks. *Anesthesiology*. 2009 Jan;110(1):182–188. PMID 19104185
3. Chelly JE, Schilling D. Thromboprophylaxis and peripheral nerve blocks in patients undergoing joint arthroplasty. *J Arthroplasty*. 2008;23:350–354. PMID 18358371
4. Chelly JE, Szczodry DM, Neumann KJ. International normalized ratio and prothrombin time values before the removal of a lumbar plexus catheter in patients receiving warfarin after total hip replacement. *Br J Anaesth*. 2008 Aug;101(2):250–254. PMID 18515818
5. Hadzic A, ed. *Textbook of Regional Anesthesia and Acute Pain Management*. New York: McGraw-Hill; 2007.
6. Horlocker TT, Wedel DJ, Rowlingson JC, et al. Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American Society of Regional Anesthesia and Pain Medicine evidence-based guidelines. *Reg Anesth Pain Med*. 2010 Jan-Feb;35(1):64–101. PMID 20052816
7. Ilfeld BM, Enneking FK. Continuous peripheral nerve blocks at home: a review. *Anesth Analg*. 2005 Jun;100(6):1822–1833. PMID 15920221